



Department of Mathematical  
& Statistical Sciences

UNIVERSITY OF COLORADO **DENVER**



# Non-intrusive Termination of Noisy Optimization

Jeffrey Larson

University of Colorado Denver

Stefan Wild

Argonne National Laboratory

FEBRUARY 24, 2012



## Problem Setting

---

When should you terminate algorithms solving

$$\min_x f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$$

when  $f$  is

- Computationally expensive
- There is noise in the computation of  $f$



## Problem Setting

---

When should you terminate algorithms solving

$$\min_x f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$$

when  $f$  is

- Computationally expensive
- There is noise in the computation of  $f$

Practitioners typically stop the optimization when:

- A measure of criticality is satisfied (e.g., gradient norm, mesh size)
- The computational budget is satisfied (e.g., number of evaluations, wall clock time)



## Problem Setting

When should you terminate algorithms solving

$$\min_x f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$$

when  $f$  is


- Computationally expensive
- There is noise in the computation of  $f$

### Definition

This is an attempt to solve the true problem:

$$\begin{array}{ll} \min_t & \text{Computational expense}(t) \\ \text{s.t.} & \text{Acceptable accuracy of the solution}(t), \end{array}$$

## Example from Nuclear Physics





## Quotes

---

*... no set of termination criteria is suitable for all optimization problems and all methods.*

*- P. Gill, W. Murray, M. Wright (Practical Optimization)*

*... it is believed that it is impossible to choose such a convergence criterion which is effective for the most general function ... so a compromise has to be made between stopping the iterative procedure too soon and calculating  $f$  an unnecessarily large number of times.*

*M. Powell (1964)*



## Quotes

---

*... no set of termination criteria is suitable for all optimization problems and all methods.*

*- P. Gill, W. Murray, M. Wright (Practical Optimization)*

*... it is believed that it is impossible to choose such a convergence criterion which is effective for the most general function ... so a compromise has to be made between stopping the iterative procedure too soon and calculating  $f$  an unnecessarily large number of times.*

*M. Powell (1964)*



## Modifications for Noisy Function

---

1. Functions with stochastic noise,  $\text{Var} \{f(x)\} > 0$ ,
  - UOBYQA, DIRECT, and Nelder-Mead methods have all been modified in the literature to repeatedly sample points.
  - Some adjust the maximum number of replications based on the noise level.
  - Termination was still based on traditional measures:
    - points clustered together
    - no decrease in the best function value
2. Functions with deterministic noise, (iterative methods, round-off error)
  - Kelley- proposes a restart technique for Nelder-Mead when low-level noise is present, but terminates independent of the noise.
  - Gramacy et al. - stops a treed Gaussian process when the maximum improvement statistic is sufficiently small.
  - Neumaier et al. - Suggest stopping SNOBFIT when the best point has not changed for a number of consecutive iterates.





## Modifications for Noisy Function

---

1. Functions with stochastic noise,  $\text{Var} \{f(x)\} > 0$ ,
  - UOBYQA, DIRECT, and Nelder-Mead methods have all been modified in the literature to repeatedly sample points.
  - Some adjust the maximum number of replications based on the noise level.
  - Termination was still based on traditional measures:
    - points clustered together
    - no decrease in the best function value
2. Functions with deterministic noise, (iterative methods, round-off error)
  - Kelley- proposes a restart technique for Nelder-Mead when low-level noise is present, but terminates independent of the noise.
  - Gramacy et al. - stops a treed Gaussian process when the maximum improvement statistic is sufficiently small.
  - Neumaier et al. - Suggest stopping SNOBFIT when the best point has not changed for a number of consecutive iterates.



## Modifications for Noisy Function

---

1. Functions with stochastic noise,  $\text{Var} \{f(x)\} > 0$ ,
  - UOBYQA, DIRECT, and Nelder-Mead methods have all been modified in the literature to repeatedly sample points.
  - Some adjust the maximum number of replications based on the noise level.
  - Termination was still based on traditional measures:
    - points clustered together
    - no decrease in the best function value
2. Functions with deterministic noise, (iterative methods, round-off error)
  - Kelley- proposes a restart technique for Nelder-Mead when low-level noise is present, but terminates independent of the noise.
  - Gramacy et al. - stops a treed Gaussian process when the maximum improvement statistic is sufficiently small.
  - Neumaier et al. - Suggest stopping SNOBFIT when the best point has not changed for a number of consecutive iterates.



## Desirable Test Properties

---

For a sequence of points and function values

$$\{x_1, \dots, x_m\} \subseteq \mathbb{R}^n, \{f_1, \dots, f_m\} \in \mathbb{R}, \mathcal{F}_i = \{(x_1, f_1), \dots, (x_i, f_i)\}$$

produced by a local minimization solver, it is preferable if the termination test is:

- Algorithm independent
  - Uses only the  $x_i$  and  $f_i$ .
- Shift and scale invariant in  $f$ 
  - Stops sequences  $\{f_i\}$  and  $\{\alpha f_i + \beta\}$  at the same point.

Useful notation: Let  $f_i^* = \min_{1 \leq j \leq i} \{f_j\}$  and  $x_i^*$  be the corresponding point.



## Desirable Test Properties

---

For a sequence of points and function values

$$\{x_1, \dots, x_m\} \subseteq \mathbb{R}^n, \{f_1, \dots, f_m\} \in \mathbb{R}, \mathcal{F}_i = \{(x_1, f_1), \dots, (x_i, f_i)\}$$

produced by a local minimization solver, it is preferable if the termination test is:

- Algorithm independent
  - Uses only the  $x_i$  and  $f_i$ .
- Shift and scale invariant in  $f$ 
  - Stops sequences  $\{f_i\}$  and  $\{\alpha f_i + \beta\}$  at the same point.

Useful notation: Let  $f_i^* = \min_{1 \leq j \leq i} \{f_j\}$  and  $x_i^*$  be the corresponding point.



## Estimate of the Noise Level

---

Let  $\hat{\varepsilon}_{i_r}$  be the relative noise in  $f_i$ .

For stochastic noise,  $\hat{\varepsilon}_{i_r} = \frac{\sqrt{\text{Var}\{f(x_i)\}}}{E\{|f(x_i)|\}}$ , in which case, for  $\alpha > 0$ :

$$\hat{\varepsilon}_{i_r} = \frac{\alpha \sqrt{\text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\alpha^2 \text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\text{Var}\{\alpha f(x_i)\}}}{E\{|\alpha f(x_i)|\}}.$$

$\hat{\varepsilon}_{i_r}$  is scale invariant.

$$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\} = \sqrt{\text{Var}\{f(x_i)\}} = \sqrt{\text{Var}\{f(x_i) + \beta\}}$$

$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\}$  is shift invariant.

For deterministic noise, invariance depends on the methods used to obtain  $\hat{\varepsilon}_{i_r}$  and  $\hat{\varepsilon}_{i_r}|f_i|$ . For one such method, see Moré & Wild (2011).



## Estimate of the Noise Level

Let  $\hat{\varepsilon}_{i_r}$  be the relative noise in  $f_i$ .

For stochastic noise,  $\hat{\varepsilon}_{i_r} = \frac{\sqrt{\text{Var}\{f(x_i)\}}}{E\{|f(x_i)|\}}$ , in which case, for  $\alpha > 0$ :

$$\hat{\varepsilon}_{i_r} = \frac{\alpha \sqrt{\text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\alpha^2 \text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\text{Var}\{\alpha f(x_i)\}}}{E\{|\alpha f(x_i)|\}}.$$

$\hat{\varepsilon}_{i_r}$  is scale invariant.

$$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\} = \sqrt{\text{Var}\{f(x_i)\}} = \sqrt{\text{Var}\{f(x_i) + \beta\}}$$

$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\}$  is shift invariant.

For deterministic noise, invariance depends on the methods used to obtain  $\hat{\varepsilon}_{i_r}$  and  $\hat{\varepsilon}_{i_r}|f_i|$ . For one such method, see Moré & Wild (2011).



## Estimate of the Noise Level

Let  $\hat{\varepsilon}_{i_r}$  be the relative noise in  $f_i$ .

For stochastic noise,  $\hat{\varepsilon}_{i_r} = \frac{\sqrt{\text{Var}\{f(x_i)\}}}{E\{|f(x_i)|\}}$ , in which case, for  $\alpha > 0$ :

$$\hat{\varepsilon}_{i_r} = \frac{\alpha \sqrt{\text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\alpha^2 \text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\text{Var}\{\alpha f(x_i)\}}}{E\{|\alpha f(x_i)|\}}.$$

$\hat{\varepsilon}_{i_r}$  is scale invariant.

$$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\} = \sqrt{\text{Var}\{f(x_i)\}} = \sqrt{\text{Var}\{f(x_i) + \beta\}}$$

$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\}$  is shift invariant.

For deterministic noise, invariance depends on the methods used to obtain  $\hat{\varepsilon}_{i_r}$  and  $\hat{\varepsilon}_{i_r}|f_i|$ . For one such method, see Moré & Wild (2011).



## Estimate of the Noise Level

---

Let  $\hat{\varepsilon}_{i_r}$  be the relative noise in  $f_i$ .

For stochastic noise,  $\hat{\varepsilon}_{i_r} = \frac{\sqrt{\text{Var}\{f(x_i)\}}}{E\{|f(x_i)|\}}$ , in which case, for  $\alpha > 0$ :

$$\hat{\varepsilon}_{i_r} = \frac{\alpha \sqrt{\text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\alpha^2 \text{Var}\{f(x_i)\}}}{\alpha E\{|f(x_i)|\}} = \frac{\sqrt{\text{Var}\{\alpha f(x_i)\}}}{E\{|\alpha f(x_i)|\}}.$$

$\hat{\varepsilon}_{i_r}$  is scale invariant.

$$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\} = \sqrt{\text{Var}\{f(x_i)\}} = \sqrt{\text{Var}\{f(x_i) + \beta\}}$$

$\hat{\varepsilon}_{i_r} E\{|f(x_i)|\}$  is shift invariant.

For deterministic noise, invariance depends on the methods used to obtain  $\hat{\varepsilon}_{i_r}$  and  $\hat{\varepsilon}_{i_r}|f_i|$ . For one such method, see Moré & Wild (2011).





## Tests on Function Values, $\phi_1$

---

For  $\nu_{\mathcal{F}_i}, \mu \in \mathbb{R}_+, \kappa \in \mathbb{N}$ . Let  $\hat{\varepsilon}_{i_r}$  be an estimate for the relative noise level of  $f_i$ .

$$\phi_1(\mathcal{F}_i; \nu_{\mathcal{F}_i}, \kappa, \mu) \text{ stops when } \frac{f_{i-\kappa+1}^* - f_i^*}{\kappa} \leq \mu |f_i^*| \nu_{\mathcal{F}_i}$$

- If  $\nu_{\mathcal{F}_i} = 1$ : stop when the average relative change in the best function value over the last  $\kappa$  evaluations is less than  $\mu$ . (scale invariant)
- If  $\nu_{\mathcal{F}_i} = \hat{\varepsilon}_{i_r}$ : stop when the average relative change in  $f^*$  is over the last  $\kappa$  evaluations is less than a factor of  $\mu$  times the relative noise. (shift and scale invariant)



## Tests on Function Values, $\phi_1$

---

For  $\nu_{\mathcal{F}_i}, \mu \in \mathbb{R}_+, \kappa \in \mathbb{N}$ . Let  $\hat{\varepsilon}_{i_r}$  be an estimate for the relative noise level of  $f_i$ .

$$\phi_1(\mathcal{F}_i; \nu_{\mathcal{F}_i}, \kappa, \mu) \text{ stops when } \frac{f_{i-\kappa+1}^* - f_i^*}{\kappa} \leq \mu |f_i^*| \nu_{\mathcal{F}_i}$$

- If  $\nu_{\mathcal{F}_i} = 1$ : stop when the average relative change in the best function value over the last  $\kappa$  evaluations is less than  $\mu$ . (scale invariant)
- If  $\nu_{\mathcal{F}_i} = \hat{\varepsilon}_{i_r}$ : stop when the average relative change in  $f^*$  is over the last  $\kappa$  evaluations is less than a factor of  $\mu$  times the relative noise. (shift and scale invariant)



## Tests on Function Values, $\phi_2$

---

For  $\nu_{\mathcal{F}_i}, \mu \in \mathbb{R}_+, \kappa \in \mathbb{N}$ . Let  $\hat{\varepsilon}_{i_r}$  be an estimate for the relative noise level of  $f_i$ .

$$\phi_2(\mathcal{F}_i; \nu_{\mathcal{F}_i}, \kappa, \mu) \text{ stops when } \max_{i-\kappa+1 \leq j \leq i} |f_j - f_i^*| \leq \mu |f_i^*| \nu_{\mathcal{F}_i}$$

- If  $\nu_{\mathcal{F}_i} = 1$ : stop when  $\kappa$  consecutive function values are within  $\mu$  of  $|f_i^*|$ .  
(scale invariant)
- If  $\nu_{\mathcal{F}_i} = \hat{\varepsilon}_{i_r}$ : stop when the maximum absolute change in  $f$  over the last  $\kappa$  evaluations is less than a factor of  $\mu$  times the noise level at  $f_i^*$ .  
(shift and scale invariant)



## Tests on Function Values, $\phi_2$


---

For  $\nu_{\mathcal{F}_i}, \mu \in \mathbb{R}_+, \kappa \in \mathbb{N}$ . Let  $\hat{\varepsilon}_{i_r}$  be an estimate for the relative noise level of  $f_i$ .

$$\phi_2(\mathcal{F}_i; \nu_{\mathcal{F}_i}, \kappa, \mu) \text{ stops when } \max_{i-\kappa+1 \leq j \leq i} |f_j - f_i^*| \leq \mu |f_i^*| \nu_{\mathcal{F}_i}$$

- If  $\nu_{\mathcal{F}_i} = 1$ : stop when  $\kappa$  consecutive function values are within  $\mu$  of  $|f_i^*|$ .  
(scale invariant)
- If  $\nu_{\mathcal{F}_i} = \hat{\varepsilon}_{i_r}$ : stop when the maximum absolute change in  $f$  over the last  $\kappa$  evaluations is less than a factor of  $\mu$  times the noise level at  $f_i^*$ .  
(shift and scale invariant)

## Dependence on the Noise Level





## Tests on $x$ Values

---

For  $\mu \in \mathbb{R}_+$ ,  $\kappa \in \mathbb{N}$ .

- $\phi_3(\mathcal{F}_i; \kappa, \mu)$  stops when  $\max_{i-\kappa+1 \leq j, k \leq i} \|x_j - x_k\| \leq \mu$ 
  - Stop when  $\kappa$  consecutive  $x$ -values are within a distance  $\mu$  of each other.
- $\phi_4(\mathcal{F}_i; \kappa, \mu)$  stops when  $\max_{i-\kappa+1 \leq j \leq i} \|x_j^* - x_i^*\| \leq \mu$ 
  - Stop when  $\kappa$  consecutive  $x_j^*$ -values are within a distance  $\mu$  of each other.
- These tests are only shift (scale) invariant if the procedure which generates the  $\{x_i\}$  is shift (scale) invariant.



## Tests on $x$ Values

---

For  $\mu \in \mathbb{R}_+$ ,  $\kappa \in \mathbb{N}$ .

- $\phi_3(\mathcal{F}_i; \kappa, \mu)$  stops when  $\max_{i-\kappa+1 \leq j, k \leq i} \|x_j - x_k\| \leq \mu$ 
  - Stop when  $\kappa$  consecutive  $x$ -values are within a distance  $\mu$  of each other.
- $\phi_4(\mathcal{F}_i; \kappa, \mu)$  stops when  $\max_{i-\kappa+1 \leq j \leq i} \|x_j^* - x_i^*\| \leq \mu$ 
  - Stop when  $\kappa$  consecutive  $x_i^*$ -values are within a distance  $\mu$  of each other.
- These tests are only shift (scale) invariant if the procedure which generates the  $\{x_i\}$  is shift (scale) invariant.



## Tests on $x$ Values

---

For  $\mu \in \mathbb{R}_+$ ,  $\kappa \in \mathbb{N}$ .

- $\phi_3(\mathcal{F}_i; \kappa, \mu)$  stops when  $\max_{i-\kappa+1 \leq j, k \leq i} \|x_j - x_k\| \leq \mu$ 
  - Stop when  $\kappa$  consecutive  $x$ -values are within a distance  $\mu$  of each other.
- $\phi_4(\mathcal{F}_i; \kappa, \mu)$  stops when  $\max_{i-\kappa+1 \leq j \leq i} \|x_j^* - x_i^*\| \leq \mu$ 
  - Stop when  $\kappa$  consecutive  $x_i^*$ -values are within a distance  $\mu$  of each other.
- These tests are only shift (scale) invariant if the procedure which generates the  $\{x_i\}$  is shift (scale) invariant.





## Comparison Test

---

As a point of comparison, we define the test

- $\phi_5(\mathcal{F}_i; \kappa)$  to stop after  $\kappa$  iterations

This test is trivially shift and scale invariant.



## Problem Set

---

53 problems of the form:

$$f(x) = 1 + (1 + \sigma g(x)) \sum_{i=1}^m F_i^s(x)^2,$$

- For stochastic noise

$$\text{Var} \{g(x)\} = 1$$

- For deterministic noise

$$g(x) = \xi(x)(4\xi(x)^2 - 3))$$

$$\xi(x) = 0.9 \sin(100\|x\|_1) \cos(100\|x\|_\infty) + 0.1 \cos(\|x\|_2).$$



## Problem Set

---

53 problems of the form:

$$f(x) = 1 + (1 + \sigma g(x)) \sum_{i=1}^m F_i^s(x)^2,$$

- For stochastic noise

$$\text{Var} \{g(x)\} = 1$$

- For deterministic noise

$$g(x) = \xi(x)(4\xi(x)^2 - 3))$$

$$\xi(x) = 0.9 \sin(100\|x\|_1) \cos(100\|x\|_\infty) + 0.1 \cos(\|x\|_2).$$



## Problem Set

---

We have 6 algorithms from the following classes:

1. Nelder-Mead implementations
2. Pattern search methods
3. Model-based methods
4. ... and methods which cross these classes

that we ran on all 53 problems, leaving us with 318 algorithm runs to form  $\mathcal{P}$ .

For each termination test  $t$  and  $p \in \mathcal{P}$ , let

$$l_{p,t}^*$$

be the number of function values required to satisfy  $t$  on problem  $p$ .



## Problem Set

---

We have 6 algorithms from the following classes:

1. Nelder-Mead implementations
2. Pattern search methods
3. Model-based methods
4. ... and methods which cross these classes

that we ran on all 53 problems, leaving us with 318 algorithm runs to form  $\mathcal{P}$ .

For each termination test  $t$  and  $p \in \mathcal{P}$ , let

$$l_{p,t}^*$$

be the number of function values required to satisfy  $t$  on problem  $p$ .



## Measures of Quality in a Stopping Point

---

- Accuracy: How far from the best point does the test stop?

$$\frac{f_{i_{p,t}}^* - f_{i_{\max}}^*}{f_{i_{p,t}}^*} \quad \text{if } i_{p,t}^* < i_{\max}$$

- Performance: Could the test have stopped sooner?

Given a collection of tests  $\mathcal{T}$ , what  $t \in \mathcal{T}$  stops when

$$f_{i_{p,t}}^* - f_{i_{\max}}^* \leq |f_{i_{p,t}}^*| \hat{\epsilon}_{i_r}$$

with the smallest  $i_{p,t}^*$ ?



## Measures of Quality in a Stopping Point

---

- Accuracy: How far from the best point does the test stop?

$$\frac{f_{i_{p,t}^*}^* - f_{i_{\max}}^*}{f_{i_{p,t}^*}^*} \quad \text{if } i_{p,t}^* < i_{\max}$$


- Performance: Could the test have stopped sooner?

Given a collection of tests  $\mathcal{T}$ , what  $t \in \mathcal{T}$  stops when

$$f_{i_{p,t}^*}^* - f_{i_{\max}}^* \leq |f_{i_{p,t}^*}^*| \hat{\varepsilon}_{i_r}$$

with the smallest  $i_{p,t}^*$ ?


## Accuracy Profiles



$$\phi_1(\mathcal{F}_i; \sigma, \kappa, \mu) \text{ stops when } \frac{f_{i-\kappa+1}^* - f_i^*}{\kappa} \leq \mu |f_i^*| \sigma$$




## Accuracy Profiles



$$\phi_1(\mathcal{F}_i; \sigma, \kappa, \mu) \text{ stops when } \frac{f_{i-\kappa+1}^* - f_i^*}{\kappa} \leq \mu |f_i^*| \sigma$$

## Performance Profiles



$\phi_1(\mathcal{F}_i; \sigma, \kappa, \mu)$  stops when  $\frac{f_{i-\kappa+1}^* - f_i^*}{\kappa} \leq \mu |f_i^*| \sigma$



## Other test

---

We performed similar analysis on the other families of tests and found the best:

- $\phi_1(\cdot, \cdot, 20n, 10^{-2})$  Stop when

$$\frac{f_{i-\kappa+1}^* - f_i^*}{20n} \leq 0.01 |f_i^*| \hat{\varepsilon}_{i_r}$$

- $\phi_2(\cdot, \cdot, 10n, 10)$  Stop when

$$\max_{i-10n+1 \leq j \leq i} |f_j - f_i^*| \leq 10 |f_i^*| \hat{\varepsilon}_{i_r}$$

- $\phi_3(\cdot, n, 10^{-7})$  Stop when

$$\max_{i-n+1 \leq j, k \leq i} \|x_j - x_k\| \leq 10^{-7}$$

- $\phi_4(\cdot, 20n, 10^{-7})$  Stop when

$$\max_{i-20n+1 \leq j \leq i} \|x_j^* - x_i^*\| \leq 10^{-7}$$



## Other test

---

We performed similar analysis on the other families of tests and found the best:

- $\phi_1(\cdot, \cdot, 20n, 10^{-2})$  Stop when

$$\frac{f_{i-\kappa+1}^* - f_i^*}{20n} \leq 0.01 |f_i^*| \hat{\varepsilon}_{i_r}$$

- $\phi_2(\cdot, \cdot, 10n, 10)$  Stop when

$$\max_{i-10n+1 \leq j \leq i} |f_j - f_i^*| \leq 10 |f_i^*| \hat{\varepsilon}_{i_r}$$

- $\phi_3(\cdot, n, 10^{-7})$  Stop when

$$\max_{i-n+1 \leq j, k \leq i} \|x_j - x_k\| \leq 10^{-7}$$

- $\phi_4(\cdot, 20n, 10^{-7})$  Stop when

$$\max_{i-20n+1 \leq j \leq i} \|x_j^* - x_i^*\| \leq 10^{-7}$$



## Other test

---

We performed similar analysis on the other families of tests and found the best:

- $\phi_1(\cdot, \cdot, 20n, 10^{-2})$  Stop when

$$\frac{f_{i-\kappa+1}^* - f_i^*}{20n} \leq 0.01 |f_i^*| \hat{\varepsilon}_{i_r}$$

- $\phi_2(\cdot, \cdot, 10n, 10)$  Stop when

$$\max_{i-10n+1 \leq j \leq i} |f_j - f_i^*| \leq 10 |f_i^*| \hat{\varepsilon}_{i_r}$$

- $\phi_3(\cdot, n, 10^{-7})$  Stop when

$$\max_{i-n+1 \leq j, k \leq i} \|x_j - x_k\| \leq 10^{-7}$$

- $\phi_4(\cdot, 20n, 10^{-7})$  Stop when

$$\max_{i-20n+1 \leq j \leq i} \|x_j^* - x_i^*\| \leq 10^{-7}$$



## Other test

We performed similar analysis on the other families of tests and found the best:

- $\phi_1(\cdot, \cdot, 20n, 10^{-2})$  Stop when

$$\frac{f_{i-\kappa+1}^* - f_i^*}{20n} \leq 0.01 |f_i^*| \hat{\varepsilon}_{i_r}$$

- $\phi_2(\cdot, \cdot, 10n, 10)$  Stop when

$$\max_{i-10n+1 \leq j \leq i} |f_j - f_i^*| \leq 10 |f_i^*| \hat{\varepsilon}_{i_r}$$


- $\phi_3(\cdot, n, 10^{-7})$  Stop when

$$\max_{i-n+1 \leq j, k \leq i} \|x_j - x_k\| \leq 10^{-7}$$

- $\phi_4(\cdot, 20n, 10^{-7})$  Stop when


$$\max_{i-20n+1 \leq j \leq i} \|x_j^* - x_i^*\| \leq 10^{-7}$$

## Most Accurate Tests






## Most Accurate Tests






## Performance Profiles for Most Accurate Tests





## Performance Profiles for Most Accurate Tests





## Recommendations for termination tests

---

Test	$\kappa$	$\mu$	Interpretation of Stopping Rule
$\phi_1$	$\approx 20n$	$\approx 0.01$	Stop when the average relative decrease in the best function value over the last $20n$ function evaluations is less than one-hundredth of the relative noise level.
$\phi_2$	$\approx 10n$	$\approx 10$	Stop when the last $10n$ function evaluations are within 10 times the absolute noise level of the best function value.
$\phi_3$	$\approx n$	$\approx 10^{-7}$	Stop when the last $n$ points evaluated are within a distance of $10^{-7}$ of each other.
$\phi_4$	$\approx 20n$	$\approx 10^{-7}$	Stop when the best point hasn't moved more a distance of $10^{-7}$ for $20n$ evaluations.



## Final Comments:

---

- Tests using knowledge of the noise are better, especially as the noise level increases.
- It is likely a better use of a computational budget to restart a stagnant algorithm.
- Nothing in these tests prevent their inclusion in
  - Derivative-based algorithms
  - The refinement stage of global algorithms
- For further information, see:

<http://www.mcs.anl.gov/~wild/tnoise>